arXiv:0808.2793v1 [physics.geo-ph] 20 Aug 2008

The Weibull - log Weibull transition of interoccurrence times for synthetic and natural earthquakes

Tomohiro Hasumi[1],* Chien-chih Chen[2],† Takuma Akimoto[1],‡ and Yoji Aizawa[1]§
[1] *Department of Applied Physics, Advanced School of Science and Engineering, Waseda University, Tokyo, Japan*
[2] *Department of Earth Sciences and Graduate Institute of Geophysics, National Central University, Jhongli, Taiwan*

(Dated: August 20, 2008)

We have studied interoccurrence time distributions by analyzing the synthetic and three natural catalogs of the Japan Meteorological Agency (JMA), the Southern California Earthquake Data Center (SCEDC), and Taiwan Central Weather Bureau (TCWB) and revealed the universal feature of the interoccurrence time statistics, Weibull - log Weibull transition. This transition reinforces the view that the interoccurrence time statistics possess Weibull statistics and log-Weibull statistics. Here in this paper, the crossover magnitude from the superposition regime to the Weibull regime m_c^2 is proportional to the plate velocity. In addition, we have found the region-independent relation, $m_c^2/m_{max} = 0.54 \pm 0.004$.

PACS numbers: 91.30.Dk, 91.30.Px, 05.65.+b, 05.45.Tp

I. INTRODUCTION

Statistical properties of time intervals between successive earthquakes, henceforward the interoccurrence times and the recurrence times, have been frequently studied in order to predict when the next big earthquake will happen. Previous papers have been focused on the determination of the probability distribution and the presentation of the scaling law, as shown in the works of [1, 2, 3, 4, 5, 6]. For instance, the Weibull distribution [1], the exponential distribution [2], the Brownian passage time (BPT) distribution [3], the generalized gamma distribution [4, 5, 6], and the log normal distribution [3] are candidates for the distribution function of interoccurrence times and recurrence times. Also, in the stationary regime, a unified scaling law was proposed by Corral [5] and then obtained by analyzing the California aftershock data [6].

Meanwhile, the time interval statistics have been studied by numerical simulations of earthquake models [7], because real earthquake data are limited. For example, both the conceptual spring-block models [8, 9] and the sophisticated Virtual California model [10] show the Weibull distribution of the recurrence times. One of the authors (TH) also reported that the survivor function of interoccurrence times in the 2D spring-block model can be described by the Zipf-Mandelbrot type power law [11], which has been observed by Abe and Suzuki [12].

Very recently, a statistical feature of the interoccurrence times, the Weibull - log Weibull transition, was proposed

---

*Electronic address: t-hasumi.1981@toki.waseda.jp
†Electronic address: chencc@earth.ncu.edu.tw
‡Electronic address: akimoto@aoni.waseda.jp
§Electronic address: aizawa@waseda.jp

TABLE I: List of Earthquake Catalogs

Catalog Name	Covering Region	Term	Number of Earthquakes	m_{min}	m_c^0
JMA [14]	25°–50°N and 125°–150°E	01/01/2001–31/10/2007	130244	2.0	2.0
SCEDC [17]	32°–37°N and 114°–122°W	01/01/2001–31/12/2007	10838	0.0	2.0
CWB [18]	21°N–26°N and 119°–123°E	01/01/2001–31/12/2007	148155	0.0	2.0
Synthetic	50 × 50 (system size)	–	297040	0.0	0.3

by analyzing the Japan Meteorological Agency (JMA) catalog [13]. We found that the probability distribution of interoccurrence times can be very well fitted by the superposition of the log-Weibull distribution and the Weibull distribution. In particular, the distribution of large earthquakes obeys the Weibull distribution with exponent less than unity indicating that the process of large earthquakes is long-term correlated. Our earlier results emphasize that the interoccurrence time statistics basically contain both Weibull and log-Weibull statistics, and as the threshold of magnitude m_c is increased, the predominant distribution could change from the log-Weibull distribution to the Weibull distribution. Those statistical properties, including the Weibull - log Weibull transition, can be also found in synthetic catalogs produced by the 2D spring-block model [8]. However, the applicability to other tectonic regions of the Weibull - log Weibull transition remains unsolved.

In this study, we further investigate the interoccurrence time statistics by analyzing the Southern California and Taiwan earthquake catalogs. Together with previous results from the JMA and synthetic catalogs, we show the universal Weibull - log Weibull transition obtained in all of the catalogs. We also demonstrate that a crossover magnitude, m_c^2, between the superposition regime and the pure Weibull regime is proportional to the plate velocity, and at the end of this paper we elucidate its implication in the geophysical sense.

II. DATA AND METHODOLOGY

For studying the interoccurrence time statistics we analyzed the JMA [14], the Southern California Earthquake Data Center (SCEDC) [17], the Taiwan Central Weather Bureau (TCWB) [18], the synthetic catalogs. The information on each catalog is listed in Table I. m_{min} and m_c^0 correspond to the minimum magnitude and the minimum cutoff magnitude, respectively. We basically consider real earthquakes with magnitude greater than and equal to 2.0 because smaller earthquakes are strongly incomplete.

The synthetic catalog is created by the 2D spring-block model with the velocity-weakening friction law [15]. This model is characterized by five parameters: the stiffness l_x^2 and l_y^2, the decrement of the friction force α, the plate velocity ν, and the difference between the maximum friction force and dynamical friction force. We set those parameters at $l_x^2 = 1, l_y^2 = 3, \alpha = 3.5, \nu = 0.01$, and $\sigma = 0.01$ for reproducing the realistic Gutenberg-Richter law [11]. Event magnitude m in the model is defined as $m = m_0 + \log_{10}\left(\sum_{i,j}^{n} \delta u_{i,j}\right)/1.5$, where $\delta u_{i,j}$ and n are the total slip at (i,j) and the number of slipping blocks, respectively. m_0 is set at 0.7, so as to shift m to positive. The occurrence time of an event is the time when a block slips for the first time during an event.

The nth interoccurrence time τ_n is defined as $t_{n+1} - t_n$, where t_n and t_{n+1} are the occurrence times of the nth and $n+1$th earthquakes, respectively. The methodology is the same as that used in our previous studies [8, 13].

TABLE II: The interoccurrence time statistics on Southern California and Taiwan. Note that the rms value calculated from the cumulative distribution of $P(\tau)$ is on order of $[10^{-3}]$. $(\cdot)$ stands for the ratio of P_w to $P(\tau)$.

		$P_w + P_{lw}$	$P_w + P_{pow}$	$P_w + P_{gam}$	$P_w + P_{ln}$
Region	m_c	rms $(P_w/P(\tau))$	rms $(P_w/P(\tau))$	rms $(P_w/P(\tau))$	rms $(P_w/P(\tau))$
California	2.0	1.7 (0.45 ± 0.006)	7.1 (0.44 ± 0.02)	25 (0.97 ± 0.05)	15 (0)
	2.5	2.3 (0.58 ± 0.007)	7.2 (0.72 ± 0.01)	17 (1)	6.0 (0)
	3.0	5.3 (0.79 ± 0.02)	6.8 (0.91 ± 0.01)	9.1 (1)	5.4 (0.57 ± 0.03)
	3.5	7.6 (1)	7.6 (1)	7.6 (1)	7.6 (1)
	4.0	23 (1)	23 (1)	23 (0.34 ± 1.29)	23 (1)
Taiwan	2.5	14 (0.40 ± 0.06)	14 (0.20 ± 0.05)	43 (0.96 ± 0.07)	32 (0)
	3.0	3.4 (0.49 ± 0.02)	7.5 (0.53 ± 0.02)	24 (0.94 ± 0.06)	12 (0)
	3.5	5.0 (0.64 ± 0.02)	8.4 (0.74 ± 0.02)	13 (0)	6.1 (0)
	4.0	3.5 (0.68 ± 0.02)	7.4 (0.89 ± 0.01)	9.3 (0.83 ± 0.03)	3.4 (0.40 ± 0.03)
	4.5	5.8 (1)	5.8 (1)	5.8 (1)	5.8 (1)
	5.0	12 (1)	12 (1)	12 (1)	12 (1)

Earthquakes with magnitude m above a threshold magnitude m_c in a studied region were considered. Then, we calculated the interoccurrence times and regressed the interoccurrence times data in the time domain of $\tau > h$. It should be noted that the spatial division was carried out when we examined JMA data. h are set at 0.5 (days) and 0 for real and synthetic earthquakes, respectively.

Our previous works [8, 13] revealed that the probability distribution of interoccurrence times $P(\tau)$ can be described by the superposition of the Weibull distribution P_w and the log-Weibull distribution P_{lw}, namely,

$$P(\tau) = p \times P_w + (1 - p) \times P_{lw} \tag{1}$$

where

$$P_w(\tau) = \left(\frac{\tau}{\tau_c}\right)^{\gamma-1} \frac{\gamma}{\tau_c} \exp\left[-\left(\frac{\tau}{\tau_c}\right)^{\gamma}\right], \; P_{lw}(\tau) = \frac{(\log(\tau/k))^{\delta-1}}{(\log\tau_c')^{\delta}} \frac{\delta}{\tau} \exp\left[-\left(\frac{\log(\tau/k)}{\log\tau_c'}\right)^{\delta}\right],$$

$\gamma, \tau_c, \delta, \tau_c'$, and k are constants. p is the ratio of P_w divided by $P(\tau)$. Obviously, $P(\tau)$ is P_{lw} when $p = 0$ and P_w when $p = 1$. k is fixed at 0.5. The log-Weibull distribution is constructed by the log modification of the cumulative distribution of the Weibull distribution. Unity δ reduces the log-Weibull distribution to the power law distribution.

III. RESULTS

The interoccurrence time statistics with different m_c for the SCEDC and TCWB data are studied. We use the Weibull distribution P_w, log-Weibull distribution P_{lw}, power law distribution P_{pow}, gamma distribution P_{gam}, log normal distribution P_{ln} and their superposition to fit the distribution of the calculated interoccurrence time data. As shown in Table. II, the distribution with larger m_c fairly follows the Weibull distribution. Comparing the obtained rms value (please see explanation in the supplementary), we found that $P(\tau)$ is best fitted by the superposition of the



FIG. 1: The cumulative distribution of interoccurrence times for different catalogs. (a), (b), and (c) correspond to the log-Weibull regime, the superposition regime, and Weibull regime, respectively. For (b) and (c), the result of the synthetic data are demonstrated in the inset figures.

TABLE III: The results of fitting parameters for different distribution functions.

caption	catalog	m_c	m_{max}	γ	τ_c	δ	τ_c'	p
(a)	Synthetic[a]	0.7	2.8	–	–	5.64± 0.01	51.3± 0.06	0
(b)	Synthetic[a]	1.6	2.8	1.15± 0.005	311± 0.73	7.36± 0.04	604± 2.04	0.69±0.05
(b)	Japan[b]	2.5	7.2	2.91± 0.14	0.79± 0.01	1.10± 0.03	1.43± 0.01	0.31±0.02
(b)	California	2.0	5.7	1.83± 0.02	0.81± 0.006	1.13± 0.01	1.48± 0.006	0.45±0.006
(b)	Taiwan	2.5	7.1	3.35± 0.55	0.60± 0.03	1.01± 0.09	1.18 ± 0.02	0.40±0.06
(c)	Synthetic[a]	1.6	2.8	1.29± 0.008	1115± 3.05	–	–	1
(c)	Japan[b]	4.5	7.2	0.82 ± 0.009	7.69± 0.07	–	–	1
(c)	California	3.5	5.7	0.95± 0.005	9.46 ± 0.04	–	–	1
(c)	Taiwan	4.5	7.1	0.92± 0.005	5.44± 0.02	–	–	1

[a] Dynamical parameters are set at $l_x^2 = 1, l_y^2 = 3, \alpha = 3.5, \nu = 0.01$, and $\sigma = 0.01$.
[b] We analyzed the data in the region, ranging from 30°N to 35°N, and ranging from 140°E to 145°E.

Weibull and log-Weibull distributions. The ratio of P_w to $P(\tau)$, p, increases as m_c is increased. Thus, we can claim that a transition from the Weibull to the log-Weibull transition occurs in both earthquake catalogs of Southern California and Taiwan. As an example, together with our previous results for the JMA and synthetic catalogs [8, 13], we show the cumulative distributions of the interoccurrence times exhibiting the log-Weibull regime, the superposition regime, and the Weibull regime in Figs. 1 (a), (b), and (c), respectively. Note that Japan data (×) in Fig. 1 is obtained from a divided region spanning 35°–40°N and 140°–145°E. In addition, the pure log-Weibull regime can only be extracted from the synthetic data as shown in Fig. 1 (a).

The maximum magnitudes, denoted by m_{max}, for the fitting results of $P(\tau)$ in Fig. 1 are listed in Table. III. For each studied area γ gradually decreases as m_c is increased. In the Weibull regime, γ obtained from real earthquakes is less than unity, whereas γ for our synthetic catalog is greater than unity. However, we have noted that γ derived from the synthetic data could be also less than 1 by tuning the parameters of the spring-block model. For all catalogs, as m_c is increased, τ_c, δ and τ_c' increases double exponentially, linearly and exponentially, respectively. Particularly,



FIG. 2: The relation between p and the scaled magnitude m_c^2/m_{max}. As for the synthetic data, the Weibull - log Weibull transition appears clearly.

p gradually increases when m_c is increased, indicating that the probability distribution changes as m_c is varied. The interoccurrence time statistics exhibit the Weibull - log Weibull transition, which means that the dominant distribution changes from the log-Weibull distribution to the Weibull distribution with increase in m_c. We thus define crossover magnitudes m_c^1 and m_c^2 representing magnitudes that the distribution change from the pure log-Weibull regime to the superposition regime and from the superposition regime to the pure Weibull regime, respectively. We focus this paper on the values of m_c^2 and they are 1.7, 4.0, 3.3, and 4.4 for the synthetic, Japan, California, and Taiwan catalogs, respectively.

Most importantly, the universality of the Weibull - log Weibull transition can be clearly shown when we consider a scaled magnitude with m_c/m_{max}. Fig. 2 demonstrates the relation between p and m_c/m_{max}, the scaled magnitude, for four catalogs. As shown in Fig. 2 p gradually increases with increases in m_c/m_{max}, and the scaled crossover magnitude m_c^2/m_{max} of the Weibull - log Weibull transition is estimated to be approximately 0.60 for all of four catalogs.

To further discuss the feature of this transition, we summarize our results obtained from 17 different regions (14 regions in Japan and other 3 regions of Southern California, Taiwan, and the synthetic data) in Fig. 3. We surprisingly find an area-independent constant for the scaled crossover magnitude, namely $m_c^2/m_{max} = 0.54 \pm 0.004$. Note that, in Fig. 3, there are three outliers whose values of m_c^2/m_{max} are less than 0.5. For those three regions of 30°–35°N and 125°–130°E, and 45°–50°N and 140°–145°E, the total numbers of earthquakes are 1739 and 1135, respectively, which are smaller than the earthquake numbers used in other regions (see Fig. 1 in Ref. [13]). Therefore, the small values of m_c^2/m_{max} are probably caused by the statistical issue of insufficient samplings. As for the region of 30°–35°N and 125°–130°E, m_{max} is 8.0 which is the largest magnitude throughout the JMA catalog we analyzed. Therefore, the scaled crossover magnitude for this region tends to be biased small.



FIG. 3: The relation between the scaled crossover magnitude m_c^2/m_{max} and the maximum magnitude m_{max} for different regions. m_c^2 ranges from 0.39 (45°–50°N and 140°–145°E) to 0.62 (Taiwan).

TABLE IV: List of the cross-over magnitude and the plate velocity (see in Ref [20, 21]). The notation of PH, EU, PA, and NA represent PHilippine sea plate, EUrasian plate, PAcific plate, and North American plate, respectively.

Region	relative plate motion	velocity [mm/yr]	m_c^2
Taiwan	PH-EU	71	4.40
East Japan	PA-PH	49	3.80 [a]
West Japan	PH-EU	47	3.78 [b]
California	PA-NA	47	3.40

[a] We take an average using three regions; 25°–30°N and 140°–145°E ($m_c^2 = 3.7$), 30°–35°N and 140°–145°E ($m_c^2 = 3.7$), and 35°–40°N and 140°–145°E ($m_c^2 = 4.0$).

[b] We take an average using five regions; 25°–30°N and 125°–130°E ($m_c^2 = 3.7$), 25°–30°N and 130°–135°E ($m_c^2 = 3.3$), 30°–35°N and 130°–135°E ($m_c^2 = 4.3$), 30°–35°N and 135°–140°E ($m_c^2 = 4.0$), and 35°–40°N and 135°–140°E ($m_c^2 = 3.6$).

IV. CONCLUSION AND DISCUSSION

We have studied the interoccurrence time statistics of natural and synthetic earthquakes by analyzing the JMA, SCEDC, TCWB, and synthetic catalogs and found the universal Weibull - log Weibull transition of the interoccurrence distribution. We emphasize that interoccurrence time statistics contain both Weibull statistics and log-Weibull statistics. And, in this paper, we demonstrate the transition does occur for different tectonic settings. We also observe the area-independent scaling relation, namely $m_c^2/m_{max} = 0.54 \pm 0.004$. Still the origin of the log-Weibull distribution and the Weibull - log Weibull transition remains open. Our present work represents the first step to fully understand the interoccurrence time statistics and the Weibull - log-Weibull transition for real earthquakes.

Although the scaled crossover magnitudes m_c^2/m_{max} is area-independent, the crossover magnitude m_c^2 from the superposition regime to the pure Weibull regime depends on the tectonic region. The geophysical implication for the area-dependent m_c^2 could be exposed when comparing the plate velocity with *averaged* m_c^2. As clearly seen from Table IV, m_c^2 is on the average proportional to the plate velocity. That means the magnitude of the largest earthquake, m_{max}, for a tectonic region is more or less proportional to the plate velocity since $m_c^2/0.54 = m_{max}$. Ruff and Kanamori showed a relation stating that the magnitude of characteristic earthquake which occurs in subduction-zone is proportional directly to the convergence rate [19], which is thus consistent with our results. This study therefore suggests a possible physical interpretation for earlier observation about the velocity-dependence of the characteristic earthquake size.

Acknowledgments

We would like to thank the JMA, SCEDC, and TCWB for allowing us to use the earthquake data. The effort of the Taiwan Central Weather Bureau to maintain the CWB Seismic Network is highly appreciated. TH is supported by the Japan Society for the Promotion of Science (JSPS) and the Earthquake Research Institute cooperative research program at the University of Tokyo. CCC is also grateful for research supports from the National Science Council (ROC) and the Department of Earth Sciences at National Central University (ROC).

---

[1] Abaimov, S. G., Turcotte, D. L., Shcherbakov, R, Rundle, J. B., Yakovlev, G, Goltz, C, and Newman, W. I (2008), Earthquakes: Recurrence and Interoccurrence Times, *Pure. Applied. Geophys., 165*, 777-795.

[2] Enescu, B., Struzik, Z., and Kiyoto, K (2008), On the recurrence time of earthquakes: insight from Vrancea (Romania) intermediate-depth events *Geophys. J. Int, 172*, 395-404.

[3] Matthews, M. V., W. L. Ellsworth, and A. P. Reasenberg (2002), A Brownian Model. for recurrent earthquakes, *Bull. Seism. Soc. Am., 92*, 2233-2250.

[4] Bak, P, Christensen, K, Danon, L, and Scanlon, T (2002), Unified Scaling Law for Earthquakes, *Phys. Rev. Lett., 88*, 178501.

[5] Corral, A (2004), Long-Term Clustering, Scaling, and Universality in the Temporal Occurrence of Earthquakes, *Phys. Rev. Lett., 92*, 108501.

[6] Shcherbakov, R, Yakovlev, G, Turcotte, D. L, and Rundle, J. B (2005), Model for the Distribution of Aftershock Interoccurrence Times, *Phys. Rev. Lett., 95*, 218501.

[7] Rundle, J. B, Turcotte, D. L., Klein, W (Eds) (2000), *GeoComplexity and the physics of Earthquakes*, AGU, Washington D. C.

[8] T. Hasumi, T. Akimoto, and Y. Aizawa, arXiv.0808.0616.

[9] Abaimov, S. G, Turcotte, D. L., Shcherbakov, R., and Rundle, J. B. (2007), Recurrence and interoccurrence behavior of self-organized complex phenomena, *Nonlin. Processes Geophys., 14*, 455-464.

[10] Yakovlev, G, Turcotte, D. L., Rundle, J. B., and Rundle, P. B. (2006), Simulation-Based Distributions of Earthquake Recurrence Times on the San Andreas Fault System, *Bull. Seismol. Soc. Am., 96(6)*, 1995-2007.

[11] Hasumi, T (2007), Interoccurrence time statistics in the two-dimensional Burridge-Knopoff earthquake model, *Phys. Rev. E, 76*, 026117.

[12] Abe, S and Suzuki, N (2005), Scale-free statistics of time interval between successive earthquakes, *Physica A, 350* 588-596.

[13] T. Hasumi, T. Akimoto, and Y. Aizawa, arXiv.0807.0485.

[14] Japan Meteorological Agency Earthquake Catalog: http://wwweic.eri.u-tokyo.ac.jp/db/jma1.

[15] Carlson, J. M., Langer, J. S., Shaw, B. E., and Tang, C (1991) Intrinsic properties of a Burridge-Knopoff model of an Earthquake, *Phys. Rev. A, 44*, 884-897.

[16] Gutenberg, B. and Richter, C. F. (1954) *Seismicity of Earth and Associated Phenomena*, 2nd ed., 310pp., Princeton Univ. Press, Princeton, N. J.

[17] Southern California Earthquake Data Center: http://www.data.scec.org/.

[18] Taiwan Central Weather Bureau: http://www.cwb.gov.tw/V5e/

[19] Ruff, L., Kanamori, H (1980), Seismicity and the subduction process. *Phys. Earth Plant. Inter., 23* 240-252.

[20] Seno, T and Alice, E. G (1993), A model for the motion of the Philippine Sea plate consistent with NUVEL-1 and Geological Data, J. Geophys. Res. 98, 17941-17948.

[21] Fowler, C. M. R. (1990), *The Solid Earth: An Introduction to Global Geophysics*, Cambridge University Press, New York.